

Shake well before use

DESCRIPTION

Ergro® Stim is a new Foliar and Seed Spray for Agricultural Crops developed with a high technology process, designed to promote the vital functions of the agricultural crops. Applied to the foliage increase the foliage development, and stimulates a better flowering and fructification, increasing quantity and quality of crops.

MAIN BENEFITS USING Ergro® Stim

- * Enriched with Fulvic Acids, that increase the penetration and transport of the L. Cistein, making it more efficient.
- * Stimulates energetically the basic vegetative growth stage, and activates the fertilizers (foliar and soil), having more assimilation.

IMPORTANT

In order to immediately confirm the benefits of **Ergro® Stim** follow your regular crop practices without **Ergro® Stim** on a small area of your field and leave it as a check plot. In another side of your field follow your regular crop practices and add **Ergro® Stim**. After 10-20 days compare your crops. You will be able to see the difference in your crop.

CAUTION


Ergro® Stim is non toxic when used as directed.

If mixed with pesticides, please follow the label directions of those products.

Do not store with food products. Do not reuse the container, destroy it. Do not leave within the reach of children.

GUARANTEE

Ergro® Stim components are guaranteed. However, as we do not have any control in its use we take no responsibility for the results obtained.



RECOMMENDATIONS AND DOSAGE

Foliar Application

Vegetables: Squash, Onion, Cauliflower, Watermelon, Melon, Potato, Tomato, Pepper, etc. Apply 6.0 fl oz per acre. Make 3 applications: at the beginning of the crop development, before flowering, and in main flowering.

Fruit Trees: Coffee, Citric, Apple, Nut, Pear, Banana, Grape, etc. Apply 6.0 fl oz per acre. Make 2 to 3 applications. 1st at flowering beginning, and the rest with 25 to 30 days between applications.

Annual Crops: Strawberry, Beans, Soy, etc. Apply 4.0-6.0 fl oz per acre. Apply at the beginning of the foliage development. Repeat after 15 days. Make 3-4 applications.

Cereals: Wheat, Corn, Sorghum, Oat, etc. Apply 4.0-6.0 fl oz per acre. Make 3-4 applications every 20 days, the first at the beginning of the spring.

It is recommended to be applied with solid foliar fertilizers, mainly with Ergro® Win and Ergro® K Excel.

Seed Application

Mix 50 cc (1.7 fl oz) of Ergro® Stim in 1-2 lt. of water, spray and cover well the amount of seeds to be used in an acre. It is recommended to be applied to the seed before planting.

Ergro® Stim a product of:



Ergo Corporación, S.A de C.V.
Maestros No. 50, Tlalnepantla,
Edo. de Mexico.
www.agroergo.com

Distributed exclusively in the United States by:



Ag Nutrition, LLC
13670 Brooks School Road
Noblesville, IN 46060
U.S.A.
Ph: (317) 288-4951
www.ag-nutrition.com

GUARANTEED ANALYSIS

Total Nitrogen (N).....	0.10%	Amoniacal Nitrogen	0.10%
Available Phosphorus (P ₂ O ₅).....	0.05%		
Soluble Potash (K ₂ O).....	0.03%		
Inorganic:			
Magnesium (Mg).....	0.26%		
Sulfur (S).....	0.79%		
Boron (B).....	0.003%		
Copper (Cu).....	0.11%		
Iron (Fe).....	0.39%		
Manganese (Mn).....	0.18%		
Molybdenum (Mo).....	0.001%		
Zinc (Zn).....	0.31%		
Organic:			
Fulvic Acid.....	5.00%		
Amino Acids.....	2.00%		
Derived from L. Cistein (Acid N-Acetil-Tiazolidin 4 Carboxilium A.A.T.C.).....	5.00%		
Folic Acid.....	0.10%		

Derived from: Zinc Sulfate, Copper Sulfate, Ferrous Sulfate, Magnesium Sulfate, Manganese Sulfate, Sodium Molybdate, Sodium Tetraboratorm (Borax), Cobalt Sulfate, Citric Acid, L. Cistein.

Fulvic Acids may enhance soil micronutrients availability within the soil, and may enhance nutrient uptake.

Caution: The application of fertilizers containing molybdenum may result in a forage crops containing levels of molybdenum which are toxic to ruminant animals.

FOR ALL CROPS



Net Content: 1 Quart (32 fl oz, .945 lt)
Net Weight: 2.5 lb (1.13 kg)



MADE IN MEXICO