

Shake well before use

## DESCRIPTION

**Ergro® Crop** is a new Foliar Spray for Agricultural Crops developed with a high technology process, added with fulvic acids, and recognized plant nutrients, essential for use in vegetal nutrition. **Ergro® Crop** contains macro and micronutrients essential for a good plant nutrition. **Ergro® Crop** is recommended for all agricultural crops effective to stimulate and promote a harmonic plant growth.

## MAIN BENEFITS USING Ergro® Crop

- \* Excellent plant aerial development.
- \* Promotes more vigorous plants.
- \* Increases quality and crop yield.

## IMPORTANT


In order to immediately confirm the benefits of **Ergro® Crop** follow your regular crop practices without **Ergro® Crop** on a small area of your field and leave it as a check plot. In another side of your field follow your regular crop practices and add **Ergro® Crop**. After 10-20 days compare your crops. You will be able to see the difference in your crop.

## CAUTION

**Ergro® Crop** is non toxic when used as directed.  
If mixed with pesticides, please follow the label directions of those products.  
Do not store with food products. Do not reuse the container, destroy it. Do not leave within the reach of children.

## GUARANTEE

**Ergro® Crop** components are guaranteed. However, as we do not have any control in its use we take no responsibility for the results obtained.



## RECOMMENDATIONS AND DOSAGE

### Foliar Application

**Vegetables:** Squash, Onion, Cauliflower, Watermelon, Melon, Potato, Tomato, Pepper, etc. Apply 6.0 fl oz per acre. Make 3 applications: at the beginning of the crop development, before flowering, and in main flowering.

**Fruit Trees:** Coffee, Citric, Apple, Nut, Pear, Banana, Grape, etc. Apply 6.0 fl oz per acre. Make 2 to 3 applications. 1<sup>st</sup> at flowering beginning, and the rest with 25 to 30 days between applications.

**Annual Crops:** Strawberry, Beans, Soy, etc. Apply 4.0-6.0 fl oz per acre. Apply at the beginning of the foliage development. Repeat after 15 days. Make 3-4 applications.

**Cereals:** Wheat, Corn, Sorghum, Oat, etc. Apply 4.0-6.0 fl oz per acre. Make 3-4 applications every 20 days, the first at the beginning of the spring.

It is recommended to be applied with solid foliar fertilizers, mainly with Ergro® Win and Ergro® K Excel.

### APPLICATION

**Ergro® Crop** is specifically designed for foliar spray applications for crops, vegetables, lawns and ornamentals, fruit and nut trees. The optimal mixture for foliar applications is one (1) part of Ergro® Crop to one hundred (100) parts of water.

**Ergro® Crop** a product of:



Ergo Corporación, S.A de C.V.  
Maestros No. 50, Tlalpanantla,  
Edo. de Mexico.

[www.agroergo.com](http://www.agroergo.com)

Distributed exclusively in the United States by:



Ag Nutrition, LLC  
13670 Brooks School Road  
Noblesville, IN 46060  
U.S.A.

Ph: (317) 288-4951  
[www.ag-nutrition.com](http://www.ag-nutrition.com)

## GUARANTEED ANALYSIS

Total Nitrogen (N).....	2.00%
0.40% Ammoniacal Nitrogen	
1.60% Urea Nitrogen	
Available Phosphorus (P <sub>2</sub> O <sub>5</sub> ).....	1.00%
Soluble Potash (K <sub>2</sub> O).....	0.60%
Total Magnesium (Mg).....	0.93%
0.93% Water Soluble Magnesium (Mg)	
Combined Sulfur (S).....	4.00%
Boron (B).....	0.012%
Cobalt (Co).....	0.001%
Water Soluble Copper (Cu).....	1.00%
Water Soluble Iron (Fe).....	1.30%
Water Soluble Manganese (Mn).....	1.10%
Molybdenum (Mo).....	0.004%
Water Soluble Zinc (Zn).....	2.30%

**Derived from:** Urea, Ammonium Sulfate, Monoammonium Phosphate, Phosphoric Acid, Potassium Sulfate, Zinc Sulfate, Copper Sulfate, Ferrous Sulfate, Magnesium Sulfate, Manganese Sulfate, Sodium Molybdate, Sodium Tetraborate, Cobalt Sulfate./F002156

### Non-Plant Food Ingredients:

Fulvic Acid.....	3.00%
Amino Acids.....	0.37%

**Derived From:** Vegetable Matter, Soy Protein.

**Fulvic Acids** may enhance soil micronutrients availability within the soil, and may enhance nutrient uptake.

**Caution:** The application of fertilizers containing molybdenum may result in a forage crops containing levels of molybdenum which are toxic to ruminant animals.

## FOR ALL CROPS



MADE IN MEXICO

Net Content: 1 Quart (32 fl oz, .945 lt)